

The Role of Gut Microbiota in Metabolism and Immune Response: A Literature Review on Metabolic Health

Delia Yusfarani¹, Uswatun Hasanah², Anita Restu Puji Raharjeng¹, Andi Saputra¹, Muhammad Alif Saputra¹, Ikhwan Hafizh Mauludi³, Kenny Edward Yap⁴

¹ Department Biology, Faculty of Science and Technology, Universitas Islam Negeri Raden Fatah Palembang, South Sumatera, Indonesia

² Department of Hadith Studies, Faculty of Ushuluddin and Philosophy Universitas Islam Negeri Raden Fatah Palembang, South Sumatera, Indonesia

³ Ar-Rasyid Islamic Hospital Palembang, South Sumatera, Indonesia

⁴ Philippine Society for Cosmetic Surgery Makati City, Metro Manila, Philippines

*Email: deliayusfarani_uin@radenfatah.ac.id

Received: 19/11/2024

Revised: 27/01/2025

Accepted: 08/02/2025

Online: 01/05/2025

ABSTRACT

The gut microbiota, a community of microorganisms within the digestive tract, plays a pivotal role in human health by regulating both immune function and metabolism. This review examines the contributions of the gut microbiota to immune responses and metabolic health, as well as its role in preventing chronic diseases. We employed a mixed-methods (qualitative and quantitative) literature review spanning the last 50 years and analyzed 100 articles related to Gut Microbiota in Metabolism and Immune Response. The inclusion criteria encompassed all studies focusing on the gut microbiota's impact on metabolic and immune functions, while non-relevant studies were excluded. Our analysis highlights recent findings on the microbiota's ability to produce metabolites such as short-chain fatty acids (SCFAs), which exhibit anti-inflammatory properties and reinforce mucosal integrity. A balanced gut microbiota supports immune regulation, inhibits pathogen colonization, and optimizes metabolic processes. However, lifestyle factors including high-sugar, low-fiber diets, antibiotic misuse, stress, and insufficient sleep can disrupt this microbial equilibrium and lead to dysbiosis. Such imbalances can precipitate metabolic and inflammatory disorders, including diabetes, autoimmune diseases, and cardiovascular conditions. This review underscores the importance of maintaining a balanced gut microbiota for disease prevention and long-term health, as well as the need to raise awareness about the impact of lifestyle choices on gut health.

Keywords: Dysbiosis; Gut Microbiota; Immune System; Metabolic Health; SCFA.

Copyright © 2025. The authors (CC BY-SA 4.0)

Introduction

The gut microbiota, composed of trillions of microorganisms within the digestive tract, plays a fundamental role in maintaining human health [1]. Since advances in next-generation sequencing and metagenomics have allowed for a deeper exploration of gut microbial diversity, it has become evident that these microbial communities are involved not

only in nutrient digestion and metabolism but also in modulating immune responses [2], [3]. Numerous studies have demonstrated that the gut microbiota produces bioactive metabolites, such as short-chain fatty acids (SCFAs), that help regulate the immune system, maintain mucosal barrier integrity, and limit inflammatory processes [2]. This interaction between the microbiota and the

host immune system is crucial for sustaining immunological homeostasis and preventing overgrowth of pathogens.

Despite of the growing body of research indicating the gut microbiota's integral part in overall health, major gaps remain. While previous studies have confirmed the gut microbiota's role in disease etiology, ranging from autoimmune disorders to metabolic syndromes, there is less clarity on how multifactorial lifestyle elements collectively influence microbiota balance over the long term. For instance, it has been established that diets high in sugar and low in fiber can reduce beneficial microbial diversity, yet the precise mechanisms by which these dietary patterns lead to chronic dysbiosis and inflammation remain partially unexplored [4], [5]. Similarly, although antibiotic misuse and inadequate sleep have been implicated in disrupting the microbiome, comprehensive investigations that integrate these factors alongside stress and extreme dietary habit are relatively scarce [6], [7].

Current evidence also points to the gut microbiota as a mediating factor in mental health via the gut-brain axis, underscoring the broader systemic implications of dysbiosis [8], [9]. However, state of the art research highlights the need for further elucidation of how specific microbial shifts contribute to metabolic and immunological dysfunction. Emerging reviews call for integrative approaches, combining metagenomic, metabolomic, and clinical data, to map the causal pathways linking gut microbial alterations to chronic diseases such as type 2 diabetes, inflammatory bowel disease, and cardiovascular disorders [10], [11].

Lifestyle behaviors, such as the consumption of fast food, high-sugar/low-fiber diets, and irregular sleep, can significantly perturb gut microbial communities, yet public awareness of these connections remains limited [12]. Under normal conditions, a balanced microbiota can outcompete pathogens via nutrient competition and immune modulation [13].

However, an imbalance (dysbiosis) often triggers chronic inflammatory states, enhances gut permeability (*leaky gut*), and increases the risk of a host of systemic diseases [14]. Recent findings indicate that dysbiosis can induce metabolic disturbances by influencing insulin resistance and oxidative stress [15]. Nevertheless, the precise interplay between gut microbial composition, immunomodulation, and metabolic pathways is still being unraveled, reinforcing the need for in-depth research in this area.

Given these critical gaps and the mounting evidence of the microbiota's far-reaching effects, this review aims to examine the mechanisms by which the gut microbiota sustains immunological balance and metabolic health, with a focus on preventing chronic diseases. By synthesizing key insights from recent and foundational studies, we seek to bridge the current knowledge gaps and shed light on targeted strategies, including dietary interventions, probiotic use, and lifestyle modifications, to uphold gut microbial homeostasis and bolster long-term well being.

Materials and Methods

Research Objectives

This study aims to analyze the role of gut microbiota and its metabolites in metabolism, immune responses within the gut, and its relationship to overall metabolic health.

Research Methods

Studies were included if they focused on the gut microbiota's role in metabolic health and immune responses, used original research designs or relevant reviews/meta-analyses, involved human or animal models with clear metabolic or immune outcomes, and were published in English within the last 50 years, contributing to a total of 100 journal articles. We excluded studies that did not directly address gut microbiota in

metabolism or immunity, lacked rigorous methods (e.g., no clear microbiota analysis), were duplicates, or had no full text available.

In this review, no meta-analysis was conducted due to the heterogeneity of study designs, populations, and outcome measures. Instead, we employed a narrative

synthesis to integrate both quantitative and qualitative findings. Additionally, we did not follow a formal systematic review protocol (e.g., PRISMA or PROSPERO); however, our methodology was structured, applying predefined inclusion and exclusion criteria, and adhering to recognized standards for literature reviews.

Table 1. Organs and Identifiable Human Microbiota [16]

Organ	Microbiota	References
Intrauterine (in the womb)	Microbiota from placental tissue, amniotic fluid, and meconium (baby's first stool): <i>Lactobacillus crispatus</i> , <i>Bifidobacterium</i> , <i>Enterococcus</i> , <i>Staphylococcus</i> , <i>Streptococcus</i> .	[17–21]
Infant gut (at birth)	Microbiota from mother's feces and vaginal microbiota (through normal delivery): <i>Bacteroides</i> , <i>Prevotella</i> , <i>Ruminococcus gnavus</i> , <i>Clostridium difficile</i> , <i>Eubacterium</i> , <i>Bifidobacterium</i> , <i>Lactobacillus</i> , <i>Enterococcus faecium</i> , <i>Escherichia coli</i> .	[22–31]
Infant gut (via cesarean birth)	Microbiota from skin and other environments: <i>Staphylococcus</i> (e.g., <i>S. epidermidis</i> , <i>Staphylococcus aureus</i>), <i>Corynebacterium</i> , <i>Propionibacterium</i> (now often classified as <i>Cutibacterium</i>), Other hospital-acquired or environmental microbes (e.g., certain strains of <i>Enterococcus</i> or <i>Enterobacter</i>).	[32–38]
Adult gut	Thousands of bacterial species, mainly from 8 of 55 primary bacterial phyla: Firmicutes (e.g., <i>Ruminococcus</i> , <i>Clostridium</i> , <i>Lactobacillus</i>), Bacteroidetes (e.g., <i>Bacteroides</i> , <i>Prevotella</i>), Actinobacteria (e.g., <i>Bifidobacterium</i>), Proteobacteria (e.g., <i>Escherichia</i> , <i>Salmonella</i>), Verrucomicrobia (e.g., <i>Akkermansia muciniphila</i>), Fusobacteria (e.g., <i>Fusobacterium</i>), Tenericutes (although often in low abundance), Euryarchaeota (archaea, like <i>Methanobrevibacter smithii</i> , sometimes considered separately from bacteria).	[39–52]
Gut epithelium	Bacteria promoting antimicrobial peptides (e.g.: α and β defensin, Reg3 γ , Ang4): Bacteroidetes (e.g., <i>Bacteroides fragilis</i>), Firmicutes (e.g., <i>Lactobacillus</i> , <i>Clostridium species</i>), <i>Akkermansia muciniphila</i> (Verrucomicrobia), <i>Enterococcus faecalis</i> (Firmicutes), <i>Faecalibacterium prausnitzii</i> (Firmicutes – Clostridiales), <i>Escherichia coli</i> (Proteobacteria – Komensal Strains), <i>Ruminococcus</i> (Firmicutes – Lachnospiraceae family).	[51], [53–60]

Results and Discussion

Results

The analysis shows (table 1 and table 2), that organ specific microbiota play

a vital role in maintaining physiological functions, while gut microbiota are essential for metabolism, nutrient absorption, and immune regulation.

Table 2. Gut Microbiota

Organ/Area	Microbiota	Main Function	References
Gut (Intestinal Tract)	<i>Faecalibacterium prausnitzii</i> , <i>Roseburia intestinalis</i> , <i>Anaerostipes butyraticus</i> , <i>Bifidobacteria</i> .	Metabolize complex carbohydrates to produce SCFAs (butyrate, acetate, propionate); regulate immune response, maintain gut barrier integrity.	[57], [61–63]
Gut Epithelium	Various gut microbiota as follows: Bacteroidetes (e.g., <i>Bacteroides fragilis</i>), Firmicutes (e.g., <i>Lactobacillus</i> , Clostridium species), Akkermansia muciniphila (Verrucomicrobia), Enterococcus faecalis (Firmicutes), <i>Faecalibacterium prausnitzii</i> (Firmicutes–Clostridiales), <i>Escherichia coli</i> (Proteobacteria–Commensal Strains), <i>Ruminococcus</i> (Firmicutes–Lachnospiraceae family).	Modulate immune response; SCFAs regulate immune cell function, epithelial cell turnover, anti-inflammatory pathways.	[51], [53–60]
Mucosal Layer (Gut Barrier)	Firmicutes, Bifidobacteriaceae.	Regulate immune tolerance, support epithelial integrity, reduce permeability to pathogens.	[64], [65]
Immune cell (Makrofag, DC)	SCFA from gut (<i>Butirat</i> , <i>Aetat</i>), the microbiota as follows: Butirat: <i>Faecalibacterium prausnitzii</i> , <i>Roseburia spp.</i> , <i>Eubacterium rectale</i> , <i>Anaerostipes spp.</i> , <i>Clostridium butyricum</i> . Aetat: <i>Bifidobacterium spp.</i> , <i>Escherichia coli</i> (non-patogen), <i>Lactobacillus spp.</i> , <i>Akkermansia muciniphila</i> . Propionat: <i>Bacteroides spp.</i> , <i>Veillonella spp.</i> , <i>Prevotella spp.</i>	SCFAs activate GPR –G Protein-coupled receptor (GPR41, GPR43, GPR109a), modulate proinflammatory cytokines, promote T-reg cell differentiation.	[57], [61, 66–72]
Colonocytes (Large Intestinal Cells)	Butyrate-producing bacteria: <i>Faecalibacterium prausnitzii</i> (Firmicutes–Clostridiales family), <i>Roseburia spp.</i> (Firmicutes – Lachnospiraceae family), <i>Eubacterium rectale</i> (Firmicutes – Lachnospiraceae family), <i>Anaerostipes spp.</i> (Firmicutes– Lachnospiraceae family), <i>Butyrivibrio spp.</i> (Firmicutes – Lachnospiraceae family), <i>Clostridium butyricum</i> (Firmicutes – Clostridiaceae family), <i>Ruminococcus spp.</i> (Firmicutes – Ruminococcaceae family).	Absorb SCFAs to support epithelial cell health and barrier function; SCFAs enhance anti-inflammatory responses, regulate cell turnover: anti-inflammatory properties by producing butyrate and modulating immune responses. Produces butyrate through the fermentation of dietary fiber. Ferments dietary fiber to produce butyrate and propionate. Plays a role in gut. Associated with improved gut barrier function and reduced inflammation. Often used in probiotic formulations for its beneficial effects on gut health and immune modulation.	[73–79]

Table 2. Continue

Organ/Area	Microbiota	Main Function	References
	Commensal anaerobic bacteria: <i>Lactobacillus</i> spp. (Firmicutes – Lactobacillaceae family), <i>Bifidobacterium</i> spp. (Actinobacteria– Bifidobacteriaceae family), <i>Clostridium</i> spp. (Firmicutes – Clostridiaceae family), <i>Prevotella</i> spp. (Bacteroidetes – Prevotellaceae family), <i>Veillonella</i> spp. (Firmicutes – Veillonellaceae family), <i>Peptostreptococcus</i> spp. (Firmicutes– Peptostreptococcaceae family), <i>Akkermansia muciniphila</i> (Verrucomicrobia family), <i>Fusobacterium</i> spp. (Fusobacteriaceae family), <i>Enterococcus</i> spp. (Firmicutes – Enterococcaceae family).	Maintain low pH, support healthy epithelial function, prevent pathogen colonization through SCFA production. Help in lactose digestion, production of lactic acid, and inhibition of pathogen colonization. Ferment complex carbohydrates and produce beneficial short-chain fatty acids (SCFAs) such as acetate. Essential energy source for intestinal epithelial cells. Associated with individuals consuming high-fiber diets. Maintaining gut pH balance and reducing the accumulation of harmful metabolites. Play a role in protein metabolism and maintaining gut homeostasis. Specializes in mucin degradation, which supports intestinal barrier integrity. Can act as opportunistic pathogens under dysbiotic conditions but are part of the normal commensal flora. Play a role in bile salt metabolism and resistance to bile acids. Major site of SCFA production, influencing immune response, inflammation regulation, and gut health. Produce acetate and lactate, which help maintain gut health. Involved in fiber digestion, SCFAs production (especially butyrate), and metabolic regulation. Degrading resistant starch and plant fibers, contributing to overall gut health. Production of butyrate: Supports colonocyte health and reduces inflammation. Fermentation of dietary fibers: Helps in breaking down plant-based fibers into SCFAs. Regulation of metabolism: Firmicutes have been linked to energy extraction from food, which can influence body weight. Immune modulation: Interact with intestinal immune cells to maintain homeostasis and prevent excessive inflammation.	[52], [56], [68], [71], [72], [80–83]
Small Intestine	<i>Bifidobacterium</i> (Phylum: Actinobacteria): <i>Bifidobacterium longum</i> , <i>Bifidobacterium breve</i> , <i>Bifidobacterium bifidum</i> , <i>Bifidobacterium adolescentis</i> . Firmicutes (Phylum: Firmicutes): Faecalibacterium prausnitzii (Order: Clostridiales), Roseburia spp. (Order: Lachnospiraceae), Clostridium spp. (Order: Clostridiaceae), Eubacterium rectale (Order: Lachnospiraceae), Blautia spp. (Order: Lachnospiraceae). Ruminococcus spp. (Order: Ruminococcaceae): Lactobacillus spp. (Order: Lactobacillales).		[57], [84–90]
Large Intestine (Colon)			

Discussion

The gut microbiota is essential in the development of human immunity (Table 1). Bacteria within the microbiota contribute to immune system maturation by modulating both innate and adaptive immune responses from an early age. Fermentation of fiber by the microbiota produces SCFAs, which act as anti-inflammatory agents, support gut epithelial health, and protect against pathogen invasion. Additionally, the microbiota regulates immune responses by stimulating antibody and immune cell production, functioning as a “shield” against pathogens [4].

Gut microbiota also impacts metabolism, producing SCFAs from fiber and complex carbohydrate fermentation, supporting neurotransmitter function, and enhancing nutrient absorption (Table 2). Short-chain fatty acids (SCFAs) are formed through the anaerobic fermentation of complex carbohydrates by gut microbes.

Acetate is produced from acetyl-CoA via acetyl phosphate. Propionate is generated through the succinate pathway or the acrylate pathway. Butyrate is produced through a series of reactions from acetyl-CoA to butyryl-CoA, which is then converted into butyrate. The main factors influencing SCFA biosynthesis include substrate availability, the composition of the microbial community, and gut environmental conditions such as pH. By understanding the pathways involved in SCFA formation, we can determine how nutritional interventions (for example, increasing fiber intake) or microbiota management can enhance the proportion of specific SCFAs that benefit health. Microbiota imbalance, or dysbiosis, can increase HPA axis activity during stress, affecting blood pressure, blood sugar levels, and other metabolic disturbances [5]. Types of Microorganisms in Gut Microbiota shows in Table 3 [6].

Table 3. Types of Microorganisms in Gut Microbiota

No	Organ	Microbiota	Roles
1	Gut Bacteria	<i>Lactobacillus</i>	Involved in fiber fermentation producing short-chain fatty acids (SCFAs)
		<i>Clostridium</i>	
		Bacteroides	Break down complex polysaccharides, playing a key role in fiber metabolism
		Actinobacteria	Participates in carbohydrate metabolism
		Proteobacteria; <i>Escherichia coli</i> Verrucomicrobi; <i>Akkermansia muciniphila</i>	Plays a metabolic role; includes aids in digestion and vitamin K synthesis
2	Gut Fungi	<i>Saccharomyces cerevisiae</i> <i>Candida spp</i>	Involved in mucus metabolism acts as a probiotic, pathogenic in dysbiosis
3	Gut Archaea	<i>Methanobrevibacter smithii</i>	involved in methane gas production from hydrogen fermentation
4	Gut Protozoa	<i>Entamoeba histolytica</i> <i>Giardia lamblia</i>	cause intestinal infections on human cause intestinal infections on human help control bacterial populations in the gut, influencing microbial interactions and potentially preventing pathogenic infections.
5	Gut Viruses	Bacteriophages	

The Importance of Gut Microbiota Balance for Metabolic Health:

An imbalance in gut microbiota can lead to inflammation and metabolic diseases such as diabetes, dyslipidemia, and hyperuricemia. Thus, maintaining a balanced microbiota is crucial for metabolic health. Various lifestyle habits can disrupt this balance, known as dysbiosis, and negatively impact gut health:

1. Processed Foods High in Sugar, Saturated Fats, and Low in Fiber

Processed foods that are high in sugar, high in saturated fats, and low in fiber are types of food that typically undergo extensive processing and the addition of various additives. This processing can include preservation, the addition of chemical substances, refining, and even modifications to texture or flavor. These foods promote the growth of pathogenic bacteria while suppressing beneficial microbes. Excessive sugar, for example, can increase pathogenic bacteria like *Clostridium* and *Escherichia coli*, potentially causing intestinal inflammation [7].

2. Antibiotics

While antibiotics kill infection-causing bacteria, they also eliminate beneficial gut microbiota, reducing bacterial diversity. Long-term or repeated antibiotic use can increase dysbiosis risk, recurrent infections, and disrupt long-term microbial [8].

3. Low Fiber Intake

Fiber is a key food source for gut microbiota, fermenting into SCFAs that are crucial for gut mucosal health and immune response. Lack of fiber can reduce SCFA-producing bacteria like *Faecalibacterium prausnitzii* and *Bifidobacteria*, which act as natural anti-inflammatories. A diet low in fiber (and often high in simple sugars and fats) frequently leads to a decrease in gut microbiota diversity. The reduction of fiber-degrading bacteria can create space for bacteria more tolerant of a high-

fat/sugar diet, such as certain species within the Firmicutes phylum [9], [10].

4. Stress

Stress elevates cortisol production, which negatively impacts gut microbiota. Cortisol increases gut permeability, allowing harmful microbes to enter the bloodstream and trigger inflammatory responses. Chronic stress may also reduce beneficial bacteria and heighten dysbiosis risk. Stress can lower the population of beneficial short-chain fatty acid (SCFA)-producing bacteria, such as *Lactobacillus*, *Bifidobacterium*, and *Faecalibacterium prausnitzii*. A reduction in these bacteria can weaken the intestinal mucosal barrier and decrease the production of crucial metabolites for gut health [5], [11].

5. Smoking and Excessive Alcohol Consumption

Smoking alters gut microbiota composition by reducing beneficial bacteria and fostering pathogenic microbes. Excessive alcohol similarly increases gut permeability, inflammation, and microbial imbalance. The chemicals in cigarette smoke (including nicotine, tar, and free radicals) can compromise the tight junctions between intestinal epithelial cells, making them “leaky.” This weakened barrier allows endotoxins and pathogenic microbes to cross into the bloodstream, increasing systemic inflammation. Alcohol and its metabolites (like acetaldehyde) also damage the gut lining. They disrupt tight junction proteins, impairing the gut’s barrier function and enabling harmful substances to leak through the intestinal wall [12], [13].

6. Insufficient Sleep or Irregular Sleep Patterns

Poor sleep disrupts the body's circadian rhythm, which also regulates gut microbiota activity. Inadequate sleep can reduce microbial diversity, increase pathogenic bacteria, and impair food metabolism [15]. The gut microbiota

follows daily (circadian) cycles in growth and activity. Poor or irregular sleep disrupts these cycles, reducing beneficial microbial populations and promoting the growth of potentially pathogenic species. Chronic sleep loss increases cortisol (a stress hormone) and disrupts melatonin and appetite-regulating hormones (like ghrelin and leptin). These imbalances alter the gut environment, potentially increasing gut permeability and inflammation, and reducing the overall resilience of beneficial microbes.

7. Lack of Physical Activity

Physical activity boosts beneficial bacteria like *Akkermansia muciniphila*, which is important for gut mucosal health and metabolic function [91,92]. Physical exercise can promote blood flow and oxygenation within the gastrointestinal tract. This enhanced circulation supports a healthier gut environment, making it more hospitable to beneficial microbes such as *Akkermansia muciniphila*. In contrast, a sedentary lifestyle may diminish these supportive factors, leading to decreased populations of beneficial bacteria.

Avoiding these behaviors can help maintain a healthy gut microbiota balance, which is vital for immune and metabolic health. A balanced gut microbiota aids in digestion, immune regulation, metabolism, and pathogen protection. Imbalance, or dysbiosis, increases the risk of multiple diseases. Below is an analysis of the importance of gut microbiota balance and preventive efforts to reduce disease risks related to dysbiosis.

a) Disease Risk Analysis from Gut Dysbiosis

Significant changes in the gut microbiota composition (not merely the appearance or disappearance of certain species) whether an increase or decrease in the proportion of specific bacterial groups can trigger or indicate dysbiosis. Essentially, many microbes are present in both healthy and dysbiotic states; however, their relative proportions shift drastically

enough to affect physiological functions and increase disease risk. An increase in Proteobacteria, opportunistic Firmicutes species, and inflammation-inducing bacteria, coupled with a decrease in *Bifidobacterium*, *Lactobacillus*, *Faecalibacterium*, *Roseburia*, *Eubacterium*, and *Akkermansia*, generally represents a dysbiotic pattern associated with various disease risks.

- 1) **Metabolic Disorders:** Dysbiosis affects body metabolism, contributing to obesity, type 2 diabetes, and metabolic syndrome. Imbalance disrupts carbohydrate and fat metabolism, increases insulin resistance and reduces cellular insulin sensitivity, heightening the risk of diabetes and obesity [93,94].
- 2) **Autoimmune and Inflammatory Diseases:** Dysbiosis impacts immune response, raising the risk of autoimmune and inflammatory diseases like inflammatory bowel disease (IBD), rheumatoid arthritis, and lupus. It can lead to increased gut permeability ("leaky gut"), allowing pathogens and toxins to enter the bloodstream and trigger excessive immune responses [94,95].
- 3) **Mental Health Disorders:** Gut microbiota communicates with the brain through the gut-brain axis. Dysbiosis may cause systemic inflammation and affect neurotransmitter production, contributing to mental health issues such as depression, anxiety, and chronic stress. [96,97].
- 4) **Cardiovascular Disease:** Certain gut microbiota metabolites can contribute to cardiovascular diseases. For instance, trimethylamine N-oxide (TMAO) from specific foods can increase atherosclerosis, hypertension, and heart disease risk. [94].

b) Prevention and Maintenance of Gut Microbiota Balance

Maintaining a balanced gut microbiota is key to preventing a range of

diseases. Some main reasons for this include:

- 1) Reduced risk of chronic diseases, especially those linked to inflammation and metabolic disorders. A balanced microbiota provides natural protection against pathogens, reducing chronic infection and inflammation risks.
- 2) A healthy microbiota enables the immune system to effectively recognize and eliminate pathogens while maintaining tolerance toward commensal organisms. This reduces the risk of autoimmune diseases.
- 3) Gut-brain axis health is supported by a balanced microbiota, helping regulate neurotransmitters like serotonin and dopamine, potentially preventing mental disorders triggered by dysbiosis.

c) Dysbiosis Prevention Strategies

Preventing gut dysbiosis involves lifestyle and dietary changes to support gut health. Recommended strategies include:

- 1) Fiber: Sources from vegetables, fruits, and whole grains feed beneficial bacteria like *Bifidobacteria* and *Faecalibacterium prausnitzii*, producing SCFAs that benefit gut health and immunity.
- 2) Antibiotic Use: Antibiotics should be used judiciously and only as prescribed by a doctor for necessary infections.
- 3) Reducing Processed Foods: Foods high in sugar and saturated fat disrupt microbiota balance by encouraging pathogen growth while reducing such foods helps maintain beneficial bacteria.
- 4) Physical Activity and Adequate Sleep: Both contribute to microbiota diversity and help reduce stress, positively impacting gut health.
- 5) Probiotics and Prebiotics: Probiotics found in yogurt or supplements increase beneficial bacteria, while prebiotics in foods like garlic, onions, and asparagus feed these microbes, supporting their growth.

Table 4. Types of Microorganisms in Human Gut

Category	Condition	Microbiota	References
Age	Gut microbiota in infants	<i>Bifidobacteria</i>	[98]
	Gut microbiota in adults	<i>Firmicutes</i>	[85]
		<i>Bacteroidetes</i>	[85]
		<i>Clostridium difficile</i>	[99]
	Gut microbiota in the elderly	<i>Bifidobacteria</i>	[100]
Residence	Urban residence	<i>Firmicutes</i>	[85]
	Rural residence	-Not mention-	[101]
Gender	Effect of sex hormones	<i>Prevotella</i>	[102]
		<i>Bifidobacteria</i>	[103]
Comorbidities	Obesity	<i>Lactobacillus</i>	[104]
		<i>Firmicutes</i>	[105]
		<i>Bacteroidetes</i>	[105]
	Type 2 diabetes	<i>Faecalibacterium prausnitzii</i>	[106]
	Autoimmune diseases	<i>Escherichia coli</i>	[107]

Certain foods can disrupt the balance of gut microbiota and reduce the number of beneficial bacteria. The widespread consumption of fast food across various groups in society highlights a low awareness of its harmful impact on gut microbiota. This consumption leads to a reduction in beneficial gut bacteria and

increases the risk of dysbiosis. Antibiotic misuse has become fairly common, with people often taking antibiotics at the slightest discomfort without medical guidance.

Most people do not consume enough fruits, vegetables, or whole grains, which are essential for supporting the

growth of beneficial gut bacteria. This trend reflects a general lack of knowledge about the role of fiber in maintaining a healthy gut microbiota. Many are also unaware that stress and poor sleep can negatively impact gut health, potentially causing digestive issues and inflammation.

Moreover, many view products that support gut health as mere trends, without understanding their role in preserving gut microbiota balance. However, few realize that certain habits affect not only organs like the lungs and liver but also disrupt the balance of bacteria in the gut. These diets are often followed without considering their impact on gut microbiota [96].

Beside that, age, residence, gender, and comorbidities are also factors that Influence the gut microbiota composition and function. Several factors, including age, residence, gender, and comorbidities, can influence the composition and function of gut microbiota. Table 4, shows types of microbiota in human gut in relation to age, residence, gender, and comorbidities, also the explanation of it.

Here is the explanation:

1. Age. The gut microbiota undergoes significant changes throughout human life, from infancy to old age

Factors such as birth mode, diet, and environmental exposure shape microbiota composition from an early age.

- Gut microbiota in infants: At birth, the infant's gut is nearly sterile. Initial colonization is influenced by the mode of delivery (vaginal or cesarean section). Vaginally delivered infants acquire a richer microbiota dominated by *Bifidobacteria* from the mother's vaginal flora, whereas cesarean-delivered infants tend to develop a microbiota resembling skin flora. Breastfeeding promotes the growth of *Bifidobacteria*, which plays a crucial role in immune system development and metabolism.
- Gut microbiota in adults: More diverse compared to infants, with dominant phyla such as *Firmicutes* and

Bacteroidetes. Diet, lifestyle, and environmental exposure play a significant role in maintaining microbiota balance.

- Gut microbiota in the elderly: Decreased microbial diversity, with an increase in opportunistic pathogens such as *Clostridium difficile*. Reduction in *Bifidobacteria* and *Firmicutes* may contribute to chronic inflammation and metabolic disorders.

2. Residence (Environment and Geography)

The living environment, such as urban or rural areas, also affects gut microbiota through diet, hygiene, and exposure to environmental microorganisms.

- Urban residence: Tends to have lower microbial diversity due to a diet high in processed foods and low in fiber. Higher antibiotic exposure and sanitation levels can reduce contact with beneficial microorganisms.
- Rural residence: Greater microbial diversity, with bacteria such as *Prevotella*, which are associated with high-fiber and complex carbohydrate diets. Greater exposure to soil and animals enhances microbial diversity.

3. Gender

Although gender differences in gut microbiota are not always prominent, research indicates that sex hormones can influence microbiota composition.

- Effect of sex hormones: Hormones such as estrogen and testosterone can impact microbial diversity and composition. Women tend to have higher levels of *Bifidobacteria* and *Lactobacillus*, which are associated with gut and reproductive health.
- Differences in disease risk related to microbiota: Women are more prone to digestive disorders such as irritable bowel syndrome (IBS), whereas men are more susceptible to metabolic disorders like microbiota-related obesity.

4. Comorbidities (Underlying Health Conditions)

Certain health conditions, such as obesity, diabetes, and autoimmune diseases, significantly affect gut microbiota composition.

- Obesity: An increase in bacteria from the *Firmicutes* phylum compared to *Bacteroidetes*, contributing to enhanced energy absorption from food. Production of inflammatory metabolites such as lipopolysaccharides (LPS), which trigger chronic inflammation.
- Type 2 diabetes: Gut microbiota associated with insulin resistance tends to have an increase in pathogenic bacteria and a decrease in SCFA (short-chain fatty acid)-producing bacteria such as *Faecalibacterium prausnitzii*.
- Autoimmune diseases (e.g., Crohn's disease and ulcerative colitis): Dysbiosis characterized by reduced microbial diversity and an increase in pro-inflammatory bacteria such as pathogenic *Escherichia coli*.

Maintaining gut microbiota balance is essential for overall health, including metabolic, immune, and mental well-being. Adopting a healthy lifestyle and avoiding risk factors for dysbiosis can significantly reduce the risk of chronic diseases associated with microbiota imbalance. Preventing dysbiosis may also extend lifespan and improve quality of life.

Conclusion

Based on the literature, gut microbiota plays a crucial role in regulating immune responses and metabolic health. Types of microorganisms in gut microbiota such as bacteria (*Lactobacillus*, *Bacteroides*, *Bifidobacterium*), fungi (*Saccharomyces cerevisiae*), archaea (*Methanobrevibacter smithii*), protozoa (*Entamoeba histolytica*), and viruses (bacteriophages) interact to influence immunity and overall health. Balanced gut microbiota is a primary factor for human health, while imbalances can trigger metabolic and immunological diseases. Although many studies demonstrate a

consistent role of the microbiota in metabolism and immune function, some findings remain contradictory or inconclusive. These discrepancies generally reflect the complex interactions between microbes and their host, various external factors (diet, lifestyle, environment), and variations in research methodologies. This situation highlights the need for more comprehensive and controlled studies, as well as a personalized approach to assessing microbiota health and implementing suitable interventions.

Conflict of interest

We declare that there is no conflict of interest.

References

- [1] Afzaal M, Saeed F, Shah YA, Hussain M, Rabail R, Socol CT, et al. Human gut microbiota in health and disease: Unveiling the relationship. *Front Microbiol* 2022;13:1–14, doi: [10.3389/fmicb.2022.999001](https://doi.org/10.3389/fmicb.2022.999001).
- [2] Morrison DJ, Preston T. Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism. *Gut Microbes* 2016;7:189–200, doi: [10.1080/19490976.2015.1134082](https://doi.org/10.1080/19490976.2015.1134082).
- [3] Gill PA, Inniss S, Kumagai T, Rahman FZ, Smith AM. The Role of Diet and Gut Microbiota in Regulating Gastrointestinal and Inflammatory Disease. *Front Immunol* 2022;13:866059, doi: [10.3389/fimmu.2022.866059](https://doi.org/10.3389/fimmu.2022.866059).
- [4] Hasibuan FEB, Kolondam dan BJ. Interaksi Antara Mikrobiota Usus Dan Sistem Kekebalan Tubuh Manusia. *J Ilm Sains* 2014;17:35, doi: [10.35799/jis.17.1.2017.15221](https://doi.org/10.35799/jis.17.1.2017.15221).
- [5] Yoo JY, Groer M, Dutra SVO, Sarkar A, McSkimming DI. Gut microbiota and immune system interactions. *Microorganisms* 2020;8:1–22, doi: [10.3390/microorganisms8101587](https://doi.org/10.3390/microorganisms8101587).

- [6] Pratama RB, Berawi KN, Islamy N. Mikrobiota Usus dan Osteoarthritis. *J Ilmu Medis Indones* 2021;1:1–6, doi: [10.35912/jimi.v1i1.279](https://doi.org/10.35912/jimi.v1i1.279).
- [7] Dinan TG, Cryan JF. Regulation of the stress response by the gut microbiota: Implications for psychoneuroendocrinology. *Psychoneuroendocrinology* 2012;37:1369–78, doi: [10.1016/j.psyneuen.2012.03.007](https://doi.org/10.1016/j.psyneuen.2012.03.007).
- [8] Huttenhower C, Gevers D, Knight R, Abubucker S, Badger JH, Chinwalla AT, et al. Structure, function and diversity of the healthy human microbiome. *Nature* 2012;486:207–14, doi: [10.1038/nature11234](https://doi.org/10.1038/nature11234).
- [9] Martínez Leo EE, Segura Campos MR. Effect of ultra-processed diet on gut microbiota and thus its role in neurodegenerative diseases. *Nutrition* 2020;71, doi: [10.1016/j.nut.2019.110609](https://doi.org/10.1016/j.nut.2019.110609).
- [10] McDonnell L, Gilkes A, Ashworth M, Rowland V, Harries TH, Armstrong D, et al. Association between antibiotics and gut microbiome dysbiosis in children: systematic review and meta-analysis. *Gut Microbes* 2021;13:1–18, doi: [10.1080/19490976.2020.1870402](https://doi.org/10.1080/19490976.2020.1870402).
- [11] Ojo O, Feng QQ, Ojo OO, Wang XH. The role of dietary fibre in modulating gut microbiota dysbiosis in patients with type 2 diabetes: A systematic review and meta-analysis of randomised controlled trials. *Nutrients* 2020;12:1–21, doi: [10.3390/nu12113239](https://doi.org/10.3390/nu12113239).
- [12] Penumutthu S, Korry BJ, Hewlett K, Belenky P. Fiber supplementation protects from antibiotic-induced gut microbiome dysbiosis by modulating gut redox potential. *Nat Commun* 2023;14:1–11, doi: [10.1038/s41467-023-40553-x](https://doi.org/10.1038/s41467-023-40553-x).
- [13] Palareti G, Legnani C, Cosmi B, Antonucci E, Erba N, Poli D, et al. Comparison between different D-Dimer cutoff values to assess the individual risk of recurrent venous thromboembolism: Analysis of results obtained in the DULCIS study. *Int J Lab Hematol* 2016;38:42–9, doi: [10.1111/ijlh.12426](https://doi.org/10.1111/ijlh.12426).
- [14] Gui X, Yang Z, Li MD. Effect of Cigarette Smoke on Gut Microbiota: State of Knowledge. *Front Physiol* 2021;12:1–14, doi: [10.3389/fphys.2021.673341](https://doi.org/10.3389/fphys.2021.673341).
- [15] Antinozzi M, Giffi M, Sini N, Gallè F, Valeriani F, De Vito C, et al. Cigarette Smoking and Human Gut Microbiota in Healthy Adults: A Systematic Review. *Biomedicines* 2022;10:1–16, doi: [10.3390/biomedicines10020510](https://doi.org/10.3390/biomedicines10020510).
- [16] Sudarmo S, Basrowi R, Chairunita C, Basrowi R. Aplikasi Klinik Probiotik Pada Bayi Dan Anak, 2018, p. 48–77.
- [17] Wu F, Kong Y, Chen W, Liang D, Xiao Q, Hu L, et al. Improvement of vaginal probiotics *Lactobacillus crispatus* on intrauterine adhesion in mice model and in clinical practice. *BMC Microbiol* 2023;23:78, doi: [10.1186/s12866-023-02823-y](https://doi.org/10.1186/s12866-023-02823-y).
- [18] Dera N, Żeber-Lubecka N, Ciebiała M, Kosińska-Kaczyńska K, Szymusik I, Massalska D, et al. Intrauterine Shaping of Fetal Microbiota. *J Clin Med* 2024;13, doi: [10.3390/jcm13175331](https://doi.org/10.3390/jcm13175331).
- [19] Li H, Fu L, Chen X, Xu H, Jing Q, Yang C, et al. Gut Microbiota and Metabolome Description of Antibiotic-Treated Neonates From Parturients With Intrauterine Infection. *Front Cell Infect Microbiol* 2022;12:817832, doi: [10.3389/fcimb.2022.817832](https://doi.org/10.3389/fcimb.2022.817832).
- [20] Wen Y, Wu Q, Zhang L, He J, Chen Y, Yang X, et al. Association of Intrauterine Microbes with Endometrial Factors in Intrauterine Adhesion Formation and after Medicine Treatment. *Pathog (Basel,*

- Switzerland) 2022;11, doi: [10.3390/pathogens11070784](https://doi.org/10.3390/pathogens11070784).
- [21] Corvaglia L, Tonti G, Martini S, Aceti A, Mazzola G, Aloisio I, et al. Influence of Intrapartum Antibiotic Prophylaxis for Group B Streptococcus on Gut Microbiota in the First Month of Life. *J Pediatr Gastroenterol Nutr* 2016;62:304–8, doi: [10.1097/MPG.0000000000000928](https://doi.org/10.1097/MPG.0000000000000928).
- [22] Sakwinska O, Foata F, Berger B, Brüssow H, Combremont S, Mercenier A, et al. Does the maternal vaginal microbiota play a role in seeding the microbiota of neonatal gut and nose? *Benef Microbes* 2017;8:763–78, doi: [10.3920/BM2017.0064](https://doi.org/10.3920/BM2017.0064).
- [23] Matharu D, Ponsero AJ, Dikareva E, Korpela K, Kolho K-L, de Vos WM, et al. Bacteroides abundance drives birth mode dependent infant gut microbiota developmental trajectories. *Front Microbiol* 2022;13:953475, doi: [10.3389/fmicb.2022.953475](https://doi.org/10.3389/fmicb.2022.953475).
- [24] Vuillermin PJ, O'Hely M, Collier F, Allen KJ, Tang MLK, Harrison LC, et al. Maternal carriage of *Prevotella* during pregnancy associates with protection against food allergy in the offspring. *Nat Commun* 2020;11:1452, doi: [10.1038/s41467-020-14552-1](https://doi.org/10.1038/s41467-020-14552-1).
- [25] Sagheddu V, Patrone V, Miragoli F, Puglisi E, Morelli L. Infant Early Gut Colonization by Lachnospiraceae: High Frequency of *Ruminococcus gnavus*. *Front Pediatr* 2016;4:57, doi: [10.3389/fped.2016.00057](https://doi.org/10.3389/fped.2016.00057).
- [26] Vasilescu I-M, Chifiriuc M-C, Pircalabioru GG, Filip R, Bolocan A, Lazăr V, et al. Gut Dysbiosis and *Clostridioides difficile* Infection in Neonates and Adults. *Front Microbiol* 2021;12:651081, doi: [10.3389/fmicb.2021.651081](https://doi.org/10.3389/fmicb.2021.651081).
- [27] Kashtanova DA, Popenko AS, Tkacheva ON, Tyakht AB, Alexeev DG, Boytsov SA. Association between the gut microbiota and diet: Fetal life, early childhood, and further life. *Nutrition* 2016;32:620–7, doi: [10.1016/j.nut.2015.12.037](https://doi.org/10.1016/j.nut.2015.12.037).
- [28] Mikami K, Kimura M, Takahashi H. Influence of maternal bifidobacteria on the development of gut bifidobacteria in infants. *Pharmaceuticals (Basel)* 2012;5:629–42, doi: [10.3390/ph5060629](https://doi.org/10.3390/ph5060629).
- [29] Zhang X, Mushajiang S, Luo B, Tian F, Ni Y, Yan W. The Composition and Concordance of *Lactobacillus* Populations of Infant Gut and the Corresponding Breast-Milk and Maternal Gut. *Front Microbiol* 2020;11:597911, doi: [10.3389/fmicb.2020.597911](https://doi.org/10.3389/fmicb.2020.597911).
- [30] Subramanya SH, Amberpet R, Chaudhary D, Nayak N, Padukone S, Bairy I, et al. Neonatal sepsis due to glycopeptide resistant *Enterococcus faecium* from colonized maternal gut- rare case evidence. *Antimicrob Resist Infect Control* 2019;8:29, doi: [10.1186/s13756-019-0490-x](https://doi.org/10.1186/s13756-019-0490-x).
- [31] Gothefors L, Carlsson B, Ahlstedt S, Hanson LA, Winberg J. Influence of maternal gut flora and colostral and cord serum antibodies on presence of *Escherichia coli* in faeces of the newborn infant. *Acta Paediatr Scand* 1976;65:225–32, doi: [10.1111/j.1651-2227.1976.tb16542.x](https://doi.org/10.1111/j.1651-2227.1976.tb16542.x).
- [32] Inchingolo F, Inchingolo AD, Palumbo I, Trilli I, Guglielmo M, Mancini A, et al. The Impact of Cesarean Section Delivery on Intestinal Microbiota: Mechanisms, Consequences, and Perspectives-A Systematic Review. *Int J Mol Sci* 2024;25, doi: [10.3390/ijms25021055](https://doi.org/10.3390/ijms25021055).
- [33] Erika L, Ingegerd A, Bill H, Robert S, Inga-Lisa S, Nils Å, et al. High Rate of Transfer of *Staphylococcus*

- aureus from Parental Skin to Infant Gut Flora. *J Clin Microbiol* 2004;42:530–4, doi: [10.1128/jcm.42.2.530-534.2004](https://doi.org/10.1128/jcm.42.2.530-534.2004).
- [34] Moles L, Gómez M, Moroder E, Bustos G, Melgar A, Del Campo R, et al. Staphylococcus epidermidis in feedings and feces of preterm neonates. *PLoS One* 2020;15:e0227823, doi: [10.1371/journal.pone.0227823](https://doi.org/10.1371/journal.pone.0227823).
- [35] Hoang DM, Levy EI, Vandenplas Y. The impact of Caesarean section on the infant gut microbiome. *Acta Paediatr* 2021;110:60–7, doi: [10.1111/apa.15501](https://doi.org/10.1111/apa.15501).
- [36] Shi Y-C, Guo H, Chen J, Sun G, Ren R-R, Guo M-Z, et al. Initial meconium microbiome in Chinese neonates delivered naturally or by cesarean section. *Sci Rep* 2018;8:3255, doi: [10.1038/s41598-018-21657-7](https://doi.org/10.1038/s41598-018-21657-7).
- [37] Rocha Martin VN, Schwab C, Krych L, Voney E, Geirnaert A, Braegger C, et al. Colonization of Cutibacterium avidum during infant gut microbiota establishment. *FEMS Microbiol Ecol* 2019;95, doi: [10.1093/femsec/fiy215](https://doi.org/10.1093/femsec/fiy215).
- [38] Stokholm J, Thorsen J, Chawes BL, Schjørring S, Krogfelt KA, Bønnelykke K, et al. Cesarean section changes neonatal gut colonization. *J Allergy Clin Immunol* 2016;138:881–889.e2, doi: [10.1016/j.jaci.2016.01.028](https://doi.org/10.1016/j.jaci.2016.01.028).
- [39] Jandhyala SM, Talukdar R, Subramanyam C, Vuyyuru H, Sasikala M, Nageshwar Reddy D. Role of the normal gut microbiota. *World J Gastroenterol* 2015;21:8787–803, doi: [10.3748/wjg.v21.i29.8787](https://doi.org/10.3748/wjg.v21.i29.8787).
- [40] Crovesy L, Masterson D, Rosado EL. Profile of the gut microbiota of adults with obesity: a systematic review. *Eur J Clin Nutr* 2020;74:1251–62, doi: [10.1038/s41430-020-0607-6](https://doi.org/10.1038/s41430-020-0607-6).
- [41] Robinson A, Wilde J, Allen-Vercoe E. Fusobacteria: physiology, form, and function, 2020, p. 95–134, doi: [10.1016/B978-0-12-819672-4.00006-4](https://doi.org/10.1016/B978-0-12-819672-4.00006-4).
- [42] Han YW. Fusobacterium nucleatum: a commensal-turned pathogen. *Curr Opin Microbiol* 2015;23:141–7, doi: [10.1016/j.mib.2014.11.013](https://doi.org/10.1016/j.mib.2014.11.013).
- [43] Park SY, Lee M, Lim SR, Kwon H, Lee YS, Kim JH, et al. Diversity and antimicrobial resistance in the streptococcus bovis/streptococcus equinus complex (Sbsec) isolated from Korean domestic ruminants. *Microorganisms* 2021;9:1–24, doi: [10.3390/microorganisms9010098](https://doi.org/10.3390/microorganisms9010098).
- [44] Gaci N, Borrel G, Tottey W, O'Toole PW, Brugère J-F. Archaea and the human gut: new beginning of an old story. *World J Gastroenterol* 2014;20:16062–78, doi: [10.3748/wjg.v20.i43.16062](https://doi.org/10.3748/wjg.v20.i43.16062).
- [45] De Filippis F, Pellegrini N, Laghi L, Gobbetti M, Ercolini D. Unusual sub-genus associations of faecal Prevotella and Bacteroides with specific dietary patterns. *Microbiome* 2016;4, doi: [10.1186/s40168-016-0202-1](https://doi.org/10.1186/s40168-016-0202-1).
- [46] Duan M, Wang Y, Zhang Q, Zou R, Guo M, Zheng H. Characteristics of gut microbiota in people with obesity. *PLoS One* 2021;16:e0255446, doi: [10.1371/journal.pone.0255446](https://doi.org/10.1371/journal.pone.0255446).
- [47] Binda C, Lopetuso LR, Rizzatti G, Gibiino G, Cennamo V, Gasbarrini A. Actinobacteria: A relevant minority for the maintenance of gut homeostasis. *Dig Liver Dis Off J Ital Soc Gastroenterol Ital Assoc Study Liver* 2018;50:421–8, doi: [10.1016/j.dld.2018.02.012](https://doi.org/10.1016/j.dld.2018.02.012).
- [48] Li J, Si H, Du H, Guo H, Dai H, Xu S, et al. Comparison of gut microbiota structure and Actinobacteria abundances in healthy young adults and elderly subjects: a pilot study. *BMC*

- Microbiol 2021;21:13, doi: [10.1186/s12866-020-02068-z](https://doi.org/10.1186/s12866-020-02068-z).
- [49] Rizzatti G, Lopetuso LR, Gibiino G, Binda C, Gasbarrini A. Proteobacteria: A Common Factor in Human Diseases. *Biomed Res Int* 2017;2017:9351507, doi: [10.1155/2017/9351507](https://doi.org/10.1155/2017/9351507).
- [50] Mukhopadhyay I, Hansen R, El-Omar EM, Hold GL. IBD-what role do Proteobacteria play? *Nat Rev Gastroenterol Hepatol* 2012;9:219–30, doi: [10.1038/nrgastro.2012.14](https://doi.org/10.1038/nrgastro.2012.14).
- [51] Geerlings SY, Kostopoulos I, de Vos WM, Belzer C. Akkermansia muciniphila in the Human Gastrointestinal Tract: When, Where, and How? *Microorganisms* 2018;6, doi: [10.3390/microorganisms6030075](https://doi.org/10.3390/microorganisms6030075).
- [52] Macchione IG, Lopetuso LR, Ianaro G, Napoli M, Gibiino G, Rizzatti G, et al. Akkermansia muciniphila: key player in metabolic and gastrointestinal disorders. *Eur Rev Med Pharmacol Sci* 2019;23:8075–83, doi: [10.26355/eurev_201909_19024](https://doi.org/10.26355/eurev_201909_19024).
- [53] Patrick S. A tale of two habitats: Bacteroides fragilis, a lethal pathogen and resident in the human gastrointestinal microbiome. *Microbiology* 2022;168, doi: [10.1099/mic.0.001156](https://doi.org/10.1099/mic.0.001156).
- [54] Wang J, Ji H, Wang S, Liu H, Zhang W, Zhang D, et al. Probiotic Lactobacillus plantarum Promotes Intestinal Barrier Function by Strengthening the Epithelium and Modulating Gut Microbiota. *Front Microbiol* 2018;9:1953, doi: [10.3389/fmicb.2018.01953](https://doi.org/10.3389/fmicb.2018.01953).
- [55] Lopetuso LR, Scaldaferri F, Petito V, Gasbarrini A. Commensal Clostridia: leading players in the maintenance of gut homeostasis. *Gut Pathog* 2013;5:23, doi: [10.1186/1757-4749-5-23](https://doi.org/10.1186/1757-4749-5-23).
- [56] Krista D, G. PE. Enterococci and Their Interactions with the Intestinal Microbiome. *Microbiol Spectr* 2017;5:10.1128/microbiolspec.bad-0014–2016, doi: [10.1128/microbiolspec.bad-0014-2016](https://doi.org/10.1128/microbiolspec.bad-0014-2016).
- [57] Miquel S, Martín R, Rossi O, Bermúdez-Humarán LG, Chatel JM, Sokol H, et al. Faecalibacterium prausnitzii and human intestinal health. *Curr Opin Microbiol* 2013;16:255–61, doi: [10.1016/j.mib.2013.06.003](https://doi.org/10.1016/j.mib.2013.06.003).
- [58] Sokol H, Seksik P, Furet JP, Firmesse O, Nion-Larmurier I, Beaugerie L, et al. Low counts of Faecalibacterium prausnitzii in colitis microbiota. *Inflamm Bowel Dis* 2009;15:1183–9, doi: [10.1002/ibd.20903](https://doi.org/10.1002/ibd.20903).
- [59] Trzeciak P, Herbet M. Role of the Intestinal Microbiome, Intestinal Barrier and Psychobiotics in Depression. *Nutrients* 2021;13, doi: [10.3390/nu13030927](https://doi.org/10.3390/nu13030927).
- [60] Vacca M, Celano G, Calabrese FM, Portincasa P, Gobbetti M, De Angelis M. The Controversial Role of Human Gut Lachnospiraceae. *Microorganisms* 2020;8, doi: [10.3390/microorganisms8040573](https://doi.org/10.3390/microorganisms8040573).
- [61] Nie K, Ma K, Luo W, Shen Z, Yang Z, Xiao M, et al. Roseburia intestinalis: A Beneficial Gut Organism From the Discoveries in Genus and Species. *Front Cell Infect Microbiol* 2021;11:757718, doi: [10.3389/fcimb.2021.757718](https://doi.org/10.3389/fcimb.2021.757718).
- [62] Kadowaki R, Tanno H, Maeno S, Endo A. Spore-forming properties and enhanced oxygen tolerance of butyrate-producing Anaerostipes spp. *Anaerobe* 2023;82:102752, doi: [10.1016/j.anaerobe.2023.102752](https://doi.org/10.1016/j.anaerobe.2023.102752).
- [63] O’Callaghan A, van Sinderen D. Bifidobacteria and Their Role as Members of the Human Gut Microbiota. *Front Microbiol* 2016;7:925, doi: [10.3389/fmicb.2016.00925](https://doi.org/10.3389/fmicb.2016.00925).
- [64] Paone P, Cani PD. Mucus barrier,

- mucins and gut microbiota: the expected slimy partners? *Gut* 2020, doi: [10.1136/gutjnl-2020-322260](https://doi.org/10.1136/gutjnl-2020-322260).
- [65] Alemao CA, Budden KF, Gomez HM, Rehman SF, Marshall JE, Shukla SD, et al. Impact of diet and the bacterial microbiome on the mucous barrier and immune disorders. *Allergy* 2021;76:714–34, doi: [10.1111/all.14548](https://doi.org/10.1111/all.14548).
- [66] Bui TPN, Mannerås-Holm L, Puschmann R, Wu H, Troise AD, Nijssse B, et al. Conversion of dietary inositol into propionate and acetate by commensal *Anaerostipes* associates with host health. *Nat Commun* 2021;12:4798, doi: [10.1038/s41467-021-25081-w](https://doi.org/10.1038/s41467-021-25081-w).
- [67] Huang T, Peng X-Y, Gao B, Wei Q-L, Xiang R, Yuan M-G, et al. The Effect of *Clostridium butyricum* on Gut Microbiota, Immune Response and Intestinal Barrier Function During the Development of Necrotic Enteritis in Chickens. *Front Microbiol* 2019;10:2309, doi: [10.3389/fmicb.2019.02309](https://doi.org/10.3389/fmicb.2019.02309).
- [68] Kaźmierczak-Siedlecka K, Roviello G, Catalano M, Polom K. Gut Microbiota Modulation in the Context of Immune-Related Aspects of *Lactobacillus* spp. and *Bifidobacterium* spp. in Gastrointestinal Cancers. *Nutrients* 2021;13, doi: [10.3390/nu13082674](https://doi.org/10.3390/nu13082674).
- [69] Ansaldo E, Slayden LC, Ching KL, Koch MA, Wolf NK, Plichta DR, et al. *Akkermansia muciniphila* induces intestinal adaptive immune responses during homeostasis. *Science* 2019;364:1179–84, doi: [10.1126/science.aaw7479](https://doi.org/10.1126/science.aaw7479).
- [70] Zafar H, Saier MHJ. Gut *Bacteroides* species in health and disease. *Gut Microbes* 2021;13:1–20, doi: [10.1080/19490976.2020.1848158](https://doi.org/10.1080/19490976.2020.1848158).
- [71] van den Bogert B, Meijerink M, Zoetendal EG, Wells JM, Kleerebezem M. Immunomodulatory properties of *Streptococcus* and *Veillonella* isolates from the human small intestine microbiota. *PLoS One* 2014;9:e114277, doi: [10.1371/journal.pone.0114277](https://doi.org/10.1371/journal.pone.0114277).
- [72] Iljazovic A, Roy U, Gálvez EJC, Lesker TR, Zhao B, Gronow A, et al. Perturbation of the gut microbiome by *Prevotella* spp. enhances host susceptibility to mucosal inflammation. *Mucosal Immunol* 2021;14:113–24, doi: [10.1038/s41385-020-0296-4](https://doi.org/10.1038/s41385-020-0296-4).
- [73] Leylabadlo HE, Ghotaslou R, Feizabadi MM, Farajnia S, Moaddab SY, Ganbarov K, et al. The critical role of *Faecalibacterium prausnitzii* in human health: An overview. *Microb Pathog* 2020;149:104344, doi: [10.1016/j.micpath.2020.104344](https://doi.org/10.1016/j.micpath.2020.104344).
- [74] Hillman ET, Kozik AJ, Hooker CA, Burnett JL, Heo Y, Kiesel VA, et al. Comparative genomics of the genus *Roseburia* reveals divergent biosynthetic pathways that may influence colonic competition among species. *Microb Genomics* 2020;6, doi: [10.1099/mgen.0.000399](https://doi.org/10.1099/mgen.0.000399).
- [75] Nilsen M, Madelen Saunders C, Leena Angell I, Arntzen MØ, Lødrup Carlsen KC, Carlsen K-H, et al. Butyrate Levels in the Transition from an Infant- to an Adult-Like Gut Microbiota Correlate with Bacterial Networks Associated with *Eubacterium Rectale* and *Ruminococcus Gnavus*. *Genes (Basel)* 2020;11, doi: [10.3390/genes11111245](https://doi.org/10.3390/genes11111245).
- [76] Allen-Vercoe E, Daigneault M, White A, Panaccione R, Duncan SH, Flint HJ, et al. *Anaerostipes hadrus* comb. nov., a dominant species within the human colonic microbiota; reclassification of *Eubacterium hadrum* Moore et al. 1976. *Anaerobe* 2012;18:523–9, doi: [10.1016/j.anaerobe.2012.09.002](https://doi.org/10.1016/j.anaerobe.2012.09.002).
- [77] Devaux CA, Million M, Raoult D.

- The Butyrogenic and Lactic Bacteria of the Gut Microbiota Determine the Outcome of Allogenic Hematopoietic Cell Transplant. *Front Microbiol* 2020;11:1642, doi: [10.3389/fmicb.2020.01642](https://doi.org/10.3389/fmicb.2020.01642).
- [78] Singh V, Lee G, Son H, Koh H, Kim ES, Unno T, et al. Butyrate producers, “The Sentinel of Gut”: Their intestinal significance with and beyond butyrate, and prospective use as microbial therapeutics. *Front Microbiol* 2023;13:1103836, doi: [10.3389/fmicb.2022.1103836](https://doi.org/10.3389/fmicb.2022.1103836).
- [79] Simpson HL, Campbell BJ. Review article: dietary fibre-microbiota interactions. *Aliment Pharmacol Ther* 2015;42:158–79, doi: [10.1111/apt.13248](https://doi.org/10.1111/apt.13248).
- [80] Million M, Tomas J, Wagner C, Lelouard H, Raoult D, Gorvel J-P. New insights in gut microbiota and mucosal immunity of the small intestine. *Hum Microbiome J* 2018;7–8:23–32, doi: [10.1016/j.humic.2018.01.004](https://doi.org/10.1016/j.humic.2018.01.004).
- [81] Grenda T, Grenda A, Domaradzki P, Krawczyk P, Kwiatek K. Probiotic Potential of *Clostridium* spp.-Advantages and Doubts. *Curr Issues Mol Biol* 2022;44:3118–30, doi: [10.3390/cimb44070215](https://doi.org/10.3390/cimb44070215).
- [82] Cheng Y, Ling Z, Li L. The Intestinal Microbiota and Colorectal Cancer. *Front Immunol* 2020;11:615056, doi: [10.3389/fimmu.2020.615056](https://doi.org/10.3389/fimmu.2020.615056).
- [83] Fukugaiti MH, Ignacio A, Fernandes MR, Ribeiro Júnior U, Nakano V, Avila-Campos MJ. High occurrence of *Fusobacterium nucleatum* and *Clostridium difficile* in the intestinal microbiota of colorectal carcinoma patients. *Brazilian J Microbiol [Publication Brazilian Soc Microbiol]* 2015;46:1135–40, doi: [10.1590/S1517-838246420140665](https://doi.org/10.1590/S1517-838246420140665).
- [84] Turrone F, Ribbera A, Foroni E, van Sinderen D, Ventura M. Human gut microbiota and bifidobacteria: from composition to functionality. *Antonie Van Leeuwenhoek* 2008;94:35–50, doi: [10.1007/s10482-008-9232-4](https://doi.org/10.1007/s10482-008-9232-4).
- [85] Mariat D, Firmesse O, Levenez F, Guimaraes V, Sokol H, Doré J, et al. The Firmicutes/Bacteroidetes ratio of the human microbiota changes with age. *BMC Microbiol* 2009;9:123, doi: [10.1186/1471-2180-9-123](https://doi.org/10.1186/1471-2180-9-123).
- [86] Tamanai-Shacoori Z, Smida I, Bousarghin L, Loreal O, Meuric V, Fong SB, et al. *Roseburia* spp.: a marker of health? *Future Microbiol* 2017;12:157–70, doi: [10.2217/fmb-2016-0130](https://doi.org/10.2217/fmb-2016-0130).
- [87] Guo P, Zhang K, Ma X, He P. *Clostridium* species as probiotics: potentials and challenges. *J Anim Sci Biotechnol* 2020;11:24, doi: [10.1186/s40104-019-0402-1](https://doi.org/10.1186/s40104-019-0402-1).
- [88] Luu TH, Michel C, Bard J-M, Dravet F, Nazih H, Bobin-Dubigeon C. Intestinal Proportion of *Blautia* sp. is Associated with Clinical Stage and Histoprognostic Grade in Patients with Early-Stage Breast Cancer. *Nutr Cancer* 2017;69:267–75, doi: [10.1080/01635581.2017.1263750](https://doi.org/10.1080/01635581.2017.1263750).
- [89] Scott KP, Duncan SH, Flint HJ. Dietary fibre and the gut microbiota. *Nutr Bull* 2008;33:201–11, doi: [10.1111/j.1467-3010.2008.00706.x](https://doi.org/10.1111/j.1467-3010.2008.00706.x).
- [90] Valeriano VD V, Balolong MP, Kang D-K. Probiotic roles of *Lactobacillus* sp. in swine: insights from gut microbiota. *J Appl Microbiol* 2017;122:554–67, doi: [10.1111/jam.13364](https://doi.org/10.1111/jam.13364).
- [91] Sun J, Fang D, Wang Z, Liu Y. Sleep Deprivation and Gut Microbiota Dysbiosis: Current Understandings and Implications. *Int J Mol Sci* 2023;24, doi: [10.3390/ijms24119603](https://doi.org/10.3390/ijms24119603).
- [92] Cataldi S, Bonavolontà V, Poli L, Clemente FM, De Candia M, Carvutto R, et al. The Relationship between Physical Activity, Physical

- Exercise, and Human Gut Microbiota in Healthy and Unhealthy Subjects: A Systematic Review. *Biology (Basel)* 2022;11, doi: [10.3390/biology11030479](https://doi.org/10.3390/biology11030479).
- [93] Dziewiecka H, Buttar HS, Kasperska A, Ostapiuk–Karolczuk J, Domagalska M, Cichoń J, et al. Physical activity induced alterations of gut microbiota in humans: a systematic review. *BMC Sports Sci Med Rehabil* 2022;14:1–22, doi: [10.1186/s13102-022-00513-2](https://doi.org/10.1186/s13102-022-00513-2).
- [94] Belizário JE, Faintuch J, Garay-Malpartida M. Gut Microbiome Dysbiosis and Immunometabolism: New Frontiers for Treatment of Metabolic Diseases. *Mediators Inflamm* 2018;2018:1–12, doi: [10.1155/2018/2037838](https://doi.org/10.1155/2018/2037838).
- [95] Amabebe E, Robert FO, Agbalalah T, Orubu ESF. Microbial dysbiosis-induced obesity: Role of gut microbiota in homeostasis of energy metabolism. *Br J Nutr* 2020;123:1127–37, doi: [10.1017/S0007114520000380](https://doi.org/10.1017/S0007114520000380).
- [96] Nibali L, Henderson B, Sadiq ST, Donos N. Insights in Chronic Inflammatory Diseases. *Microbiology* 2014;1:1–10.
- [97] Rogers GB, Keating DJ, Young RL, Wong ML, Licinio J, Wesselingh S. From gut dysbiosis to altered brain function and mental illness: Mechanisms and pathways. *Mol Psychiatry* 2016;21:738–48, doi: [10.1038/mp.2016.50](https://doi.org/10.1038/mp.2016.50).
- [98] Di Gioia D, Aloisio I, Mazzola G, Biavati B. Bifidobacteria: their impact on gut microbiota composition and their applications as probiotics in infants. *Appl Microbiol Biotechnol* 2014;98:563–77, doi: [10.1007/s00253-013-5405-9](https://doi.org/10.1007/s00253-013-5405-9).
- [99] Rea MC, O’Sullivan O, Shanahan F, O’Toole PW, Stanton C, Ross RP, et al. *Clostridium difficile* carriage in elderly subjects and associated changes in the intestinal microbiota. *J Clin Microbiol* 2012;50:867–75, doi: [10.1128/JCM.05176-11](https://doi.org/10.1128/JCM.05176-11).
- [100] Arbolea S, Watkins C, Stanton C, Ross RP. Gut Bifidobacteria Populations in Human Health and Aging. *Front Microbiol* 2016;7:1204, doi: [10.3389/fmicb.2016.01204](https://doi.org/10.3389/fmicb.2016.01204).
- [101] Saarenpää M, Roslund MI, Puhakka R, Grönroos M, Parajuli A, Hui N, et al. Do Rural Second Homes Shape Commensal Microbiota of Urban Dwellers? A Pilot Study among Urban Elderly in Finland. *Int J Environ Res Public Health* 2021;18, doi: [10.3390/ijerph18073742](https://doi.org/10.3390/ijerph18073742).
- [102] Gupta S, Khandait M, Khunger S. Exploring the gut microbiota of rural region of Haryana (India): Sociodemographic, socioeconomic factors and lifestyle. *Clin Epidemiol Glob Heal* 2024;30:101806, doi: [10.1016/j.cegh.2024.101806](https://doi.org/10.1016/j.cegh.2024.101806).
- [103] Zhang J, Sun Z, Jiang S, Bai X, Ma C, Peng Q, et al. Probiotic *Bifidobacterium lactis* V9 Regulates the Secretion of Sex Hormones in Polycystic Ovary Syndrome Patients through the Gut-Brain Axis. *MSystems* 2019;4, doi: [10.1128/mSystems.00017-19](https://doi.org/10.1128/mSystems.00017-19).
- [104] Yoon K, Kim N. Roles of Sex Hormones and Gender in the Gut Microbiota. *J Neurogastroenterol Motil* 2021;27:314–25, doi: [10.5056/jnm20208](https://doi.org/10.5056/jnm20208).
- [105] Indiani CMDSP, Rizzardi KF, Castelo PM, Ferraz LFC, Darrieux M, Parisotto TM. Childhood Obesity and Firmicutes/Bacteroidetes Ratio in the Gut Microbiota: A Systematic Review. *Child Obes* 2018;14:501–9, doi: [10.1089/chi.2018.0040](https://doi.org/10.1089/chi.2018.0040).
- [106] Remely M, Hippe B, Zanner J, Aumüller E, Brath H, Haslberger AG. Gut Microbiota of Obese, Type 2 Diabetic Individuals is Enriched in *Faecalibacterium prausnitzii*, *Akkermansia muciniphila* and

- Peptostreptococcus anaerobius after Weight Loss. Endocr Metab Immune Disord Drug Targets 2016;16:99–106, doi: [10.2174/1871530316666160831093813](https://doi.org/10.2174/1871530316666160831093813).
- [107] Lee HJ, Lee SW, Cha HR, Ha EK, Kim JH, Shin SY, et al. Acquired susceptibility to autoimmune diseases in pediatric patients with Escherichia coli infection: A population-matched retrospective cohort study. J Autoimmun 2023;137:102997, doi: [10.1016/j.jaut.2023.102997](https://doi.org/10.1016/j.jaut.2023.102997).